

Spherically-symmetric gravitational fields in the metric-affine gauge theory of gravitation

A. V. Minkevich^{1,2}, Yu. G. Vasilevski¹

¹Department of Theoretical Physics, Belarussian State University, Minsk, Belarus.

²Department of Physics and Computer Methods, University of Warmia and Mazury in Olsztyn, Poland.

E-mail: minkav@bsu.by; awm@matman.uwm.edu.pl

Abstract

Geometric structure of spherically-symmetric space-time in metric-affine gauge theory of gravity is studied. Restrictions on curvature tensor and Bianchi identities are obtained. By using certain simple gravitational Lagrangian the solution of gravitational equations for vacuum spherically-symmetric gravitational field is obtained.

gr-qc/0301098

As it is known, the application of gauge approach to gravitational interaction leads to generalization of Einsteinian theory of gravitation. At present there are different gauge theories of gravitation in dependence on using gauge group corresponding to gravitational interaction. The metric-affine gauge theory of gravitation (MAGT) is one of the most general gauge theories of gravity and it is based on the group of affine transformations $A(4, \mathbb{R})$ as gauge groupe [1]. In MAGT space-time continuum possesses curvature, torsion and nonmetricity, and as sources of gravitational field are energy-momentum tensor and so-called hypermomentum which is generalization of spin-momentum tensor of Poincare gauge theory of gravitation (PGT).

The system of gravitational equations of MAGT is complicated system of nonlinear differential equations. Their analysis is simplified in the case of models with high space symmetries. Homogeneous isotropic models in MAGT were investigated in Ref. [2]. In the framework of MAGT spherically-symmetric models are analyzed in present paper.

The geometric structure of space-time in MAGT is determined by three tensors: metrics $g_{\mu\nu}$, torsion $S^\lambda_{\mu\nu}$ and nonmetricity $Q_{\lambda\mu\nu}$ ¹. By using the system of spherical coordinates

¹ μ, ν, \dots are holonomic indices; i, k, \dots are anholonomic (tetrad) indices. Numerical tetrad indices are denoted by means of a sign $\hat{}$ over them.

($x^0 = ct$, $x^1 = r$, $x^2 = \theta$, $x^3 = \varphi$), we write metrics in the following form

$$g_{\mu\nu} = \text{diag}(e^\nu, -e^\lambda, -r^2, -r^2 \sin^2 \theta), \quad (1)$$

where $\nu = \nu(r, t)$, $\lambda = \lambda(r, t)$ are two functions of radial coordinate r and time t . The structure of tensors $S^\lambda_{\mu\nu}$ and $Q_{\lambda\mu\nu}$ in spherically-symmetric case was studied in Ref. [3]. The torsion is determined by 8 functions $S_i = S_i(r, t)$ ($i = 1, 2, \dots, 8$) and nonmetricity — by 12 functions $Q_k = Q_k(r, t)$ ($k = 0, 1, \dots, 11$). Namely nonvanishing components of tensors $S_{\lambda\mu\nu}$ and $Q_{\lambda\mu\nu}$ are:

$$\begin{aligned} S_{001} &= S_1, \quad S_{212} = S_2, \quad S_{101} = S_3, \quad S_{202} = S_4, \quad S_{313} = S_2 \sin^2 \theta, \\ S_{303} &= S_4 \sin^2 \theta, \quad S_{032} = S_5 \sin \theta, \quad S_{132} = S_6 \sin \theta, \\ S_{302} &= -S_{203} = S_7 \sin \theta, \quad S_{312} = -S_{213} = S_8 \sin \theta, \end{aligned} \quad (2)$$

$$\begin{aligned} Q_{000} &= Q_0, \quad Q_{001} = Q_1, \quad Q_{010} = Q_2, \quad Q_{011} = Q_3, \quad Q_{110} = Q_4, \\ Q_{111} &= Q_5, \quad Q_{022} = Q_6, \quad Q_{122} = Q_7, \quad Q_{220} = Q_8, \quad Q_{221} = Q_9, \\ Q_{023} &= -Q_{032} = Q_{10} \sin \theta, \quad Q_{123} = -Q_{132} = Q_{11} \sin \theta, \\ Q_{033} &= Q_6 \sin^2 \theta, \quad Q_{133} = Q_7 \sin^2 \theta, \quad Q_{330} = Q_8 \sin^2 \theta, \quad Q_{331} = Q_9 \sin^2 \theta. \end{aligned} \quad (3)$$

Note that functions S_i ($i = 5, 6, 7, 8$) and Q_k ($k = 10, 11$) have pseudoscalar character. All other components of tensors $S_{\lambda\mu\nu}$ and $Q_{\lambda\mu\nu}$ vanish, with the exception of components connected with components (2) – (3) by symmetry properties of torsion $S_{\lambda\mu\nu} = -S_{\lambda\nu\mu}$ and nonmetricity $Q_{\lambda\mu\nu} = Q_{\mu\lambda\nu}$.

By choosing diagonal tetrad h^i_μ corresponding to metrics (1)

$$h^i_\mu = \text{diag}(e^{\frac{\nu}{2}}, e^{\frac{\lambda}{2}}, r, r \sin \theta), \quad (4)$$

we find anholonomic connection:

$$A^{ik}_\mu = h^{k\nu}(\partial_\mu h^i_\nu - h^i_\lambda \Gamma^\lambda_{\nu\mu}), \quad (5)$$

where holonomic connection $\Gamma^\lambda_{\mu\nu} = \left\{ \begin{smallmatrix} \lambda \\ \mu\nu \end{smallmatrix} \right\} + S^\lambda_{\mu\nu} + S_{\mu\nu}{}^\lambda + S_{\nu\mu}{}^\lambda + \frac{1}{2}(Q_{\mu\nu}{}^\lambda - Q_\mu{}^\lambda{}_\nu - Q_\nu{}^\lambda{}_\mu)$ and $\left\{ \begin{smallmatrix} \lambda \\ \mu\nu \end{smallmatrix} \right\}$ are Christoffel symbols. Nonvanishing components of connection A^{ik}_μ are:

$$\begin{aligned} A^{\hat{0}\hat{0}}_0 &= A_0, \quad A^{\hat{0}\hat{0}}_1 = A_1, \quad A^{\hat{1}\hat{0}}_0 = A_2, \quad A^{\hat{1}\hat{0}}_1 = A_3, \quad A^{\hat{2}\hat{0}}_2 = A_4, \\ A^{\hat{3}\hat{0}}_3 &= A_4 \sin \theta, \quad A^{\hat{0}\hat{1}}_0 = A_5, \quad A^{\hat{0}\hat{1}}_1 = A_6, \quad A^{\hat{1}\hat{1}}_0 = A_7, \\ A^{\hat{1}\hat{1}}_1 &= A_8, \quad A^{\hat{2}\hat{1}}_2 = A_9, \quad A^{\hat{3}\hat{1}}_3 = A_9 \sin \theta, \quad A^{\hat{0}\hat{2}}_2 = A_{10}, \end{aligned}$$

$$\begin{aligned}
A^{\hat{0}\hat{3}}_3 &= A_{10} \sin \theta, \quad A^{\hat{1}\hat{2}}_2 = A_{11}, \quad A^{\hat{1}\hat{3}}_3 = A_{11} \sin \theta, \quad A^{\hat{2}\hat{2}}_0 = A^{\hat{3}\hat{3}}_0 = A_{12}, \\
A^{\hat{2}\hat{2}}_1 &= A^{\hat{3}\hat{3}}_1 = A_{13}, \quad A^{\hat{0}\hat{3}}_2 = A_{14}, \quad A^{\hat{0}\hat{2}}_3 = -A_{14} \sin \theta, \quad A^{\hat{1}\hat{3}}_2 = A_{15}, \\
A^{\hat{1}\hat{2}}_3 &= -A_{15} \sin \theta, \quad A^{\hat{2}\hat{3}}_0 = -A^{\hat{3}\hat{2}}_0 = A_{16}, \quad A^{\hat{2}\hat{3}}_1 = -A^{\hat{3}\hat{2}}_1 = A_{17}, \\
A^{\hat{3}\hat{0}}_2 &= A_{18}, \quad A^{\hat{2}\hat{0}}_3 = -A_{18} \sin \theta, \quad A^{\hat{3}\hat{1}}_2 = A_{19}, \quad A^{\hat{2}\hat{1}}_3 = -A_{19} \sin \theta,
\end{aligned} \tag{6}$$

where explicit form of functions A_i ($i = 0, 2, \dots, 19$) is:

$$\begin{aligned}
A_0 &= \frac{1}{2}e^{-\nu}Q_0, \quad A_1 = \frac{1}{2}e^{-\nu}Q_1, \quad A_2 = \frac{1}{2}e^{-\frac{1}{2}(\lambda+\nu)}(Q_1 - 2Q_2 + 4S_1 - e^\nu\nu'), \\
A_3 &= -\frac{1}{2}e^{-\frac{1}{2}(\lambda+\nu)}(Q_4 - 4S_3 + e^\lambda\dot{\lambda}), \quad A_4 = -\frac{1}{2r}e^{-\frac{1}{2}\nu}(Q_8 - 4S_4), \\
A_5 &= \frac{1}{2}e^{-\frac{1}{2}(\lambda+\nu)}(-Q_1 - 4S_1 + e^\nu\nu'), \quad A_6 = \frac{1}{2}e^{-\frac{1}{2}(\lambda+\nu)}(-2Q_3 + Q_4 - \\
&4S_3 + e^\lambda\dot{\lambda}), \quad A_7 = \frac{1}{2}e^{-\lambda}Q_4, \quad A_8 = \frac{1}{2}e^{-\lambda}Q_5, \quad A_9 = \frac{1}{2r}e^{-\frac{1}{2}\lambda}(2r + Q_9 - 4S_2), \\
A_{10} &= \frac{1}{2r}e^{-\frac{1}{2}\nu}(Q_8 - 2Q_6 - 4S_4), \\
A_{11} &= -\frac{1}{2r}e^{-\frac{1}{2}\lambda}(2r - 2Q_7 + Q_9 - 4S_2), \quad A_{12} = \frac{1}{2r^2}Q_8, \\
A_{13} &= \frac{1}{2r^2}Q_9, \quad A_{14} = \frac{1}{r}e^{-\frac{1}{2}\nu}S_5, \quad A_{15} = -\frac{1}{r}e^{-\frac{1}{2}\lambda}S_6, \\
A_{16} &= \frac{1}{r^2}(Q_{10} - S_5 - 2S_7), \quad A_{17} = \frac{1}{r^2}(Q_{11} - S_6 - 2S_8), \\
A_{18} &= \frac{1}{r}e^{-\frac{1}{2}\nu}(Q_{10} - S_5), \quad A_{19} = -\frac{1}{r}e^{-\frac{1}{2}\lambda}(Q_{11} - S_6).
\end{aligned} \tag{7}$$

The curvature tensor can be calculated according to his definition:

$$F^{ik}_{\mu\nu} = 2\partial_{[\mu}A^{ik}_{\nu]} + 2A^i_{l[\nu}A^{lk}_{\mu]}. \tag{8}$$

In considered case the curvature is determined by 27 functions F_i ($i = 0, 1, \dots, 26$) depending on functions ν, λ, S_i, Q_k :

$$\begin{aligned}
F^{\hat{0}\hat{0}}_{\hat{1}\hat{0}} &= F_0, \quad F^{\hat{0}\hat{1}}_{\hat{1}\hat{0}} = F_1, \quad F^{\hat{0}\hat{2}}_{\hat{2}\hat{0}} = F^{\hat{0}\hat{3}}_{\hat{3}\hat{0}} = F_2, \quad F^{\hat{0}\hat{2}}_{\hat{2}\hat{1}} = F^{\hat{0}\hat{3}}_{\hat{3}\hat{1}} = F_3, \\
F^{\hat{1}\hat{0}}_{\hat{1}\hat{0}} &= F_4, \quad F^{\hat{1}\hat{1}}_{\hat{1}\hat{0}} = F_5, \quad F^{\hat{1}\hat{2}}_{\hat{2}\hat{0}} = F^{\hat{1}\hat{3}}_{\hat{3}\hat{0}} = F_6, \quad F^{\hat{1}\hat{2}}_{\hat{2}\hat{1}} = F^{\hat{1}\hat{3}}_{\hat{3}\hat{1}} = F_7, \\
F^{\hat{2}\hat{0}}_{\hat{2}\hat{0}} &= F^{\hat{3}\hat{0}}_{\hat{3}\hat{0}} = F_8, \quad F^{\hat{2}\hat{0}}_{\hat{2}\hat{1}} = F^{\hat{3}\hat{0}}_{\hat{3}\hat{1}} = F_9, \quad F^{\hat{3}\hat{1}}_{\hat{3}\hat{0}} = F^{\hat{2}\hat{1}}_{\hat{2}\hat{0}} = F_{10},
\end{aligned}$$

$$\begin{aligned}
F^{\hat{3}\hat{1}}_{\hat{3}\hat{1}} &= F^{\hat{2}\hat{1}}_{\hat{2}\hat{1}} = F_{11}, \quad F^{\hat{2}\hat{2}}_{\hat{1}\hat{0}} = F^{\hat{3}\hat{3}}_{\hat{1}\hat{0}} = F_{12}, \quad -F^{\hat{3}\hat{2}}_{\hat{3}\hat{2}} = F^{\hat{2}\hat{3}}_{\hat{3}\hat{2}} = F_{13}, \\
F^{\hat{0}\hat{0}}_{\hat{3}\hat{2}} &= F_{14}, \quad F^{\hat{0}\hat{1}}_{\hat{3}\hat{2}} = F_{15}, \quad F^{\hat{1}\hat{0}}_{\hat{3}\hat{2}} = F_{16}, \quad F^{\hat{1}\hat{1}}_{\hat{3}\hat{2}} = F_{17}, \\
F^{\hat{2}\hat{0}}_{\hat{3}\hat{0}} &= -F^{\hat{3}\hat{0}}_{\hat{2}\hat{0}} = F_{18}, \quad F^{\hat{2}\hat{0}}_{\hat{3}\hat{1}} = -F^{\hat{3}\hat{0}}_{\hat{2}\hat{1}} = F_{19}, \quad F^{\hat{0}\hat{2}}_{\hat{3}\hat{0}} = -F^{\hat{0}\hat{3}}_{\hat{2}\hat{0}} = F_{20}, \\
F^{\hat{0}\hat{2}}_{\hat{3}\hat{1}} &= -F^{\hat{0}\hat{3}}_{\hat{2}\hat{1}} = F_{21}, \quad F^{\hat{3}\hat{1}}_{\hat{2}\hat{0}} = -F^{\hat{2}\hat{1}}_{\hat{3}\hat{0}} = F_{22}, \quad F^{\hat{3}\hat{1}}_{\hat{2}\hat{1}} = -F^{\hat{2}\hat{1}}_{\hat{3}\hat{1}} = F_{23}, \\
F^{\hat{1}\hat{2}}_{\hat{3}\hat{1}} &= -F^{\hat{1}\hat{3}}_{\hat{2}\hat{1}} = F_{24}, \quad F^{\hat{1}\hat{2}}_{\hat{3}\hat{0}} = -F^{\hat{1}\hat{3}}_{\hat{2}\hat{0}} = F_{25}, \quad F^{\hat{3}\hat{2}}_{\hat{1}\hat{0}} = -F^{\hat{2}\hat{3}}_{\hat{1}\hat{0}} = F_{26}, \\
F^{\hat{2}\hat{2}}_{\hat{3}\hat{2}} &= F^{\hat{3}\hat{3}}_{\hat{3}\hat{2}} = \frac{1}{2}(F_{14} - F_{17}).
\end{aligned} \tag{9}$$

Explicit form of functions F_i is

$$\begin{aligned}
F_0 &= \frac{1}{2}e^{-\frac{3}{2}(\lambda+\nu)}[-4Q_3S_1 - e^\lambda(\dot{Q}_1 - Q'_0 + Q_0\nu' + Q_2\dot{\lambda} - \\
&\quad Q_1\dot{\nu}) + Q_2(2Q_3 - Q_4 + 4S_3) + Q_3(e^\nu\nu' - Q_1)], \\
F_1 &= \frac{1}{4}e^{-(\lambda+2\nu)}[Q_0(Q_4 - 2Q_3 - 4S_3 + e^\lambda\dot{\lambda}) + Q_1(Q_1 + 4S_1 + e^\nu\lambda')] + \\
&\quad \frac{1}{4}e^{-(2\lambda+\nu)}\{Q_5(Q_1 + 4S_1) + Q_4^2 + e^\lambda[4\dot{Q}_3 - 2\dot{Q}_4 + 8\dot{S}_3 - 2Q'_1 - 8S'_1 + \\
&\quad 4S_1(\lambda' + \nu') - e^\nu(\lambda'\nu' + \nu'^2 + 2\nu'') - (4S_3 + 2Q_3 - Q_4)(\dot{\lambda} + \dot{\nu}) + \\
&\quad Q_4\dot{\lambda} - e^\lambda(\dot{\lambda}^2 + 2\ddot{\lambda} - \dot{\lambda}\dot{\nu})] - Q_4(2Q_3 + 4S_3) - e^\nu Q_5\nu'\}, \\
F_2 &= -\frac{1}{4r^4}e^{-(\lambda+\nu)}\{r^2(2r - 2Q_7 + Q_9 - 4S_2)(4S_1 + Q_1 - e^\nu\nu') + \\
&\quad e^\lambda[4S_5(Q_{10} - S_5 - 2S_7) - Q_8^2 + 2r^2(\dot{Q}_8 - 2\dot{Q}_6 - 4\dot{S}_4 + 2S_4\dot{\nu}) + \\
&\quad Q_8(4S_4 + 2Q_6 - r^2\dot{\nu}) + 2r^2Q_6\dot{\nu}]\} - \frac{1}{4r^2}e^{-2\nu}Q_0(2Q_6 - Q_8 + 4S_4), \\
F_3 &= -\frac{1}{4r^2}e^{-\frac{1}{2}(\lambda+\nu)}[\frac{4}{r^2}S_5(Q_{11} - S_6 + 2S_8) - 8S'_4 + 8S_4\nu' + 2(Q'_8 - 2Q'_6 + \\
&\quad 2\nu'Q_6 + \nu'Q_8)] + \frac{1}{4r^2}e^{-\frac{3}{2}\lambda-\nu}(2r + Q_9 - 2Q_7 - 4S_4)(Q_4 - 2Q_3 - 4S_3 + \\
&\quad e^\lambda\dot{\lambda}) - \frac{1}{4r^2}e^{-\frac{1}{2}\lambda-\frac{3}{2}\nu}(2Q_6 - Q_8 + 4S_4)[Q_1 + e^\nu(\frac{2}{r} + \frac{1}{r^2}Q_9 - \nu')], \\
F_4 &= \frac{1}{4}e^{-(\lambda+2\nu)}[Q_1^2 + Q_0(Q_4 - 4S_3 + e^\lambda\dot{\lambda}) + Q_1(4S_1 - 2Q_2 - e^\nu\lambda' - \\
&\quad 2e^\nu\nu')] + \frac{1}{4}e^{-(2\lambda+\nu)}\{Q_5(Q_1 + 4S_1) + Q_4^2 + e^\lambda[2\dot{Q}_4 - 8\dot{S}_3 + 4S_3(\dot{\lambda} + \dot{\nu}) + \\
&\quad e^\lambda(\dot{\lambda}^2 + 2\ddot{\lambda} - \dot{\lambda}\dot{\nu}) - Q_4\dot{\nu} + 2Q'_1 - 4Q'_2 + 8S'_1 + (2Q_2 - 4S_1)(\lambda' + \\
&\quad \nu') + e^\nu(\lambda'\nu' - \nu'^2 - 2\nu'')]\} - 2Q_2Q_5 - 4Q_4S_3 - e^\nu Q_5\nu'\},
\end{aligned} \tag{10}$$

$$\begin{aligned}
F_5 &= \frac{1}{2}e^{-\frac{3}{2}(\lambda+\nu)}[Q_3(e^\nu\nu' - Q_1 - 4S_1) + e^\nu(Q_5\dot{\lambda} - \dot{Q}_5 + Q'_4 - Q_4\lambda') + \\
&\quad Q_2(2Q_3 - Q_4 + 4S_3 - e^\lambda\dot{\lambda})], \\
F_6 &= -\frac{1}{4r^4}e^{-\frac{1}{2}(\lambda+\nu)}[(-r^2e^{-\lambda}Q_4 + Q_8 + r^2\dot{\lambda})(2r - 2Q_7 + \\
&\quad Q_9 - 4S_2) + S_6(-4Q_{10} + 4S_5 + 8S_7) + r^2(4\dot{Q}_7 - 2\dot{Q}_9 + 8\dot{S}_2) + \\
&\quad r^2(2Q_6 - Q_8 + 4S_4)(-\nu' + e^{-\nu}Q_1 + 4e^{-\nu}S_1 - 2e^{-\nu}Q_2)], \\
F_7 &= -\frac{1}{4r^4}[-e^{-2\lambda}r^2Q_5(2r - 2Q_7 + Q_9 - 4S_2) + e^{-(\lambda+\nu)}r^2(2Q_6 - Q_8 + 4S_4) \\
&\quad (4S_3 - Q_4 - e^\lambda\dot{\lambda})] - \frac{1}{4r^4}e^{-\lambda}[Q_9(4r + Q_9 - 4S_2) - 8rS_2 + 4S_6(S_6 - Q_{11} + \\
&\quad 2S_8) + 2r^2(2Q'_7 - Q'_9 + 4S'_2 + \lambda'r + \frac{1}{2}\lambda'Q_9 - 2\lambda'S_2) - 2Q_7(Q_9 + 2r + r^2\lambda')], \\
F_8 &= \frac{1}{4r^4}e^{-\nu}[(Q_8 - 4S_4)(Q_8 + e^{-\nu}r^2Q_0 - r^2\dot{\nu}) - 4(Q_{10} - S_5)(Q_{10} - S_5 - \\
&\quad 2S_7) + 2r^2(\dot{Q}_8 - 4\dot{S}_4) + e^{-\lambda}r^2(2r + Q_9 - 4S_2)(Q_1 - 2Q_2 + 4S_1 - e^\nu\nu')], \\
F_9 &= \frac{1}{4r^4}e^{-\frac{1}{2}(\lambda+\nu)}[(Q_8 - 4S_4)(Q_9 - 2r + e^{-\nu}r^2Q_1 - r^2\nu') - 4(Q_{10} - S_5)(Q_{11} - \\
&\quad S_6 - 2S_8) - e^{-\lambda}r^2(2r + Q_9 - 4S_2)(Q_4 - 4S_3 + e^\lambda\dot{\lambda}) + 2r^2(Q'_8 - 4S'_4)], \\
F_{10} &= -\frac{1}{4r^4}e^{-\frac{1}{2}(\lambda+\nu)}\{(2r + Q_9 - 4S_4)[Q_8 - e^{-\lambda}r^2(Q_4 + e^\lambda\dot{\lambda})] - 4(Q_{11} - \\
&\quad S_6)(Q_{10} - S_5 - 2S_7) + 2r^2(\dot{Q}_9 - 4\dot{S}_2) + e^{-\nu}r^2(Q_8 - 4S_4)(Q_1 + 4S_1 - e^\nu\nu')\}, \\
F_{11} &= -\frac{1}{4r^4}e^{-\lambda}[4r^2 - 4(Q_{11} - S_6)(Q_{11} - S_6 - 2S_8) + \\
&\quad e^{-\nu}r^2(Q_8 - 4S_4)(2Q_3 - Q_4 + 4S_3 - e^\lambda\dot{\lambda}) + 2r^2(Q'_9 - 4S'_2) + \\
&\quad (2r + Q_9 - 4S_2)(Q_9 - 2r - r^2e^{-\lambda}Q_5 - r^2\lambda')], \\
F_{12} &= -\frac{1}{2r^3}e^{-\frac{1}{2}(\lambda+\nu)}[2Q_8 + r(\dot{Q}_9 - Q'_8)], \\
F_{13} &= -\frac{1}{r^2} + \frac{1}{4r^4}e^{-\lambda}[(2r + Q_9 - 4S_2)(2r - 2Q_7 + Q_9 - 4S_2) + 4S_6(S_6 - \\
&\quad Q_{11})] - \frac{1}{4r^4}e^{-\nu}[(Q_8 - 4S_4)(Q_8 - 2Q_6 - 4S_4) + 4S_5(S_5 - Q_{10})], \\
F_{14} &= \frac{1}{r^4}e^{-\nu}[Q_{10}(Q_8 - 2Q_6 - 4S_4) + 2Q_6S_5], \\
F_{15} &= -\frac{1}{r^4}e^{-\frac{1}{2}(\lambda+\nu)}[(Q_{11} - S_6)(Q_8 - 2Q_6 - 4S_4) + S_5(2r + Q_9 - 4S_2)], \\
F_{16} &= -\frac{1}{r^4}e^{-\frac{1}{2}(\lambda+\nu)}[(Q_{10} - S_5)(2r - 2Q_7 + Q_9 - 4S_2) + S_6(Q_8 - 4S_4)], \\
F_{17} &= \frac{1}{r^4}e^{-\lambda}[Q_{11}(2r - 2Q_7 + Q_9 - 4S_2) + 2Q_7S_6],
\end{aligned}$$

$$\begin{aligned}
F_{18} &= \frac{1}{2r^4}e^{-\nu}\{(Q_8 - 4S_4)(Q_{10} - S_5 - 2S_7) + 2r^2(\dot{Q}_{10} - \dot{S}_5) + (Q_{10} - S_5)[Q_8 + e^{-\nu}r^2(Q_0 - e^\nu\dot{\nu})] + e^{-\lambda}r^2(Q_{11} - S_6)(Q_1 - 2Q_2 + 4S_1 - e^\nu\nu')\}, \\
F_{19} &= \frac{1}{2r^4}e^{-\frac{1}{2}(\lambda+\nu)}\{(Q_{10} - S_5)[Q_9 - 2r + e^{-\nu}r^2(Q_1 - e^\nu\nu')] + (Q_8 - 4S_4)(Q_{11} - S_6 - 2S_8) - e^{-\lambda}r^2(Q_{11} - S_6)(Q_4 - 4S_3 + e^\lambda\dot{\lambda}) + 2r^2(Q'_{10} - S'_5)\}, \\
F_{20} &= -\frac{1}{2r^4}\{e^{-2\nu}r^2Q_0S_5 - e^{-(\lambda+\nu)}r^2S_6(Q_1 + 4S_1) + e^{-\nu}[-Q_{10}(2Q_6 - Q_8 + 4S_4) - 2Q_8S_7 + (S_5 + 2S_7)(4S_4 + 2Q_6) - 2r^2\dot{S}_5 + r^2S_5\dot{\nu}] + e^{-\lambda}r^2S_6\nu'\}, \\
F_{21} &= \frac{1}{2r^4}e^{-\frac{3}{2}\lambda - \frac{1}{2}\nu}S_6[r^2(2Q_3 - Q_4 + 4S_3) + e^\lambda(Q_8 - 2Q_6 - 4S_4 - r^2\dot{\lambda})] + \frac{1}{r^4}e^{-\frac{1}{2}(\lambda+\nu)}[\frac{1}{2}Q_{11}(2Q_6 - Q_8 + 4S_4) + S_8(Q_8 - 2Q_6 - 4S_4) + r^2S'_5] - \frac{1}{2r^4}e^{-\frac{1}{2}\lambda - \frac{3}{2}\nu}S_5[r^2Q_1 + e^\nu(Q_9 + 2r + r^2\nu')], \\
F_{22} &= -\frac{1}{2r^4}e^{-\frac{1}{2}(\lambda+\nu)}[(Q_{11} - S_6)(e^{-\lambda}r^2Q_4 + r^2\dot{\lambda} - Q_8) - (2r + Q_9 - 4S_2)(Q_{10} - S_5 - 2S_7) - 2r^2(\dot{Q}_{11} - \dot{S}_6) + e^{-\nu}r^2(Q_{10} - S_5)(-Q_1 - 4S_1 + e^\nu\nu')], \\
F_{23} &= -\frac{1}{2r^4}e^{-\lambda}[(Q_{11} - S_6)(2r - Q_9 + e^{-\lambda}r^2Q_5 + r^2\lambda') - (2r + Q_9 - 4S_2)(Q_{11} - S_6 - 2S_8) - e^{-\nu}r^2(Q_{10} - S_5)(2Q_9 - Q_4 + 4S_3 - e^\lambda\dot{\lambda}) - 2r^2(Q'_{11} - S'_6)], \\
F_{24} &= \frac{1}{2r^4}e^{-2\lambda-\nu}\{e^\lambda r^2S_5(Q_4 - 4S_3) - r^2e^\nu Q_5S_6 + e^{\lambda+\nu}[2S_6(Q_7 + 2S_2) + (Q_{11} - 2S_8)(2r - 2Q_7 + Q_9 - 4S_2) + r^2S_6\lambda' - 2r^2S'_6] + e^{2\lambda}r^2S_5\dot{\lambda}\}, \\
F_{25} &= \frac{1}{2r^4}e^{-\frac{1}{2}(\lambda+\nu)}[Q_8S_6 + (2r - 2Q_7 + Q_9 - 4S_2)(Q_{10} - 2S_7 - e^\nu S_5) + r^2(S_6\dot{\lambda} - 2\dot{S}_6) + r^2S_5(2Q_2 - Q_1 - 4S_1 + e^\nu\nu')] - \frac{1}{2r^2}e^{-\frac{1}{2}(3\lambda+\nu)}Q_4S_6, \\
F_{26} &= \frac{1}{r^3}e^{-\frac{1}{2}(\lambda+\nu)}[2Q_{10} - 2S_5 - 4S_7 + r(\dot{S}_6 - 2\dot{S}_8 - Q'_{10} + S'_5 + 2S'_7)].
\end{aligned}$$

Let us consider Bianchi identities, which can be written in the following form:

$$\varepsilon^{\sigma\lambda\mu\nu}\nabla_\lambda F^i_{k\mu\nu} = 0. \quad (11)$$

where ∇ is differential operator defined analogously to covariant derivative determined by means of Cristoffel symbols or connection $(-A^{ik}_\mu)$ in case of holonomic and anholonomic indices respectively:

$\nabla_\lambda F^i_{k\mu\nu} = \partial_\lambda F^i_{k\mu\nu} + A^l_{k\lambda}F^i_{l\mu\nu} - A^i_{l\lambda}F^l_{k\mu\nu} - \left\{\begin{smallmatrix}\sigma \\ \mu\lambda\end{smallmatrix}\right\}F^i_{k\sigma\nu} - \left\{\begin{smallmatrix}\sigma \\ \nu\lambda\end{smallmatrix}\right\}F^i_{k\mu\sigma}$, and $\varepsilon^{\sigma\lambda\mu\nu}$ is Levi-Chivita symbol.

In spherically-symmetric case Bianchi identities are reduced to 20 relations:

$$\begin{aligned}
& e^{\frac{1}{2}(\lambda+\nu)}(4F_{14} + 2r^2F'_{14}) + e^\lambda[2F_{19}(2Q_6 - Q_8 + 4S_4) - \\
& 2F_{21}(Q_8 - 4S_4) + 4F_3(Q_{10} - S_5) - 4F_9S_5 + r^2\dot{\lambda}(F_{15} + F_{16})] + \\
& r^2[(F_{15} + F_{16})(Q_4 - 4S_3) - 2F_{16}Q_3] = 0, \\
& -rF_{15}(e^\lambda rQ_1 + e^\nu rQ_5) + e^{\lambda+\nu}[2F_{21}(2r + Q_9 - 4S_2) + 4rF_{15} - 4F_3(Q_{11} - \\
& S_6) + 2r^2F'_{15}] + e^{\frac{1}{2}(\lambda+\nu)}[-e^\lambda(2F_{23}(2Q_6 - Q_8 + 4S_4) + \\
& 4F_{11}S_5) + r^2(F_{17} + F_{14})(Q_4 - 2Q_3 - 4S_4 + e^\lambda\dot{\lambda})] = 0, \\
& rF_{16}(e^\lambda rQ_1 + e^\nu rQ_5) + 2e^{\lambda+\nu}[2rF_{16} + F_{19}(2r - 2Q_7 + Q_9 - 4S_2) + \\
& 2F_9S_6 + r^2F'_{16}] + e^{\frac{1}{2}(\lambda+\nu)}[r^2(F_{17} + F_{14})(Q_4 - 4S_3 + e^\lambda\dot{\lambda}) + \\
& e^\lambda F_{24}(8S_4 - 2Q_8 + 4Q_{10} - 4S_5)] = 0, \\
& 4rF_{17} + 2(F_{24} - F_{23})(2r - 2Q_7 + Q_9 - 4S_2) + 4F_{24}Q_7 - 4F_7(Q_{11} - S_6) + \\
& 4F_{11}S_6 + e^{-\frac{1}{2}(\lambda+\nu)}r^2[(F_{15} + F_{16})(Q_4 - 4S_3 + e^\lambda\dot{\lambda}) - 2F_{16}Q_3] + 2r^2F'_{17} = 0, \\
& 4r(F_{14} - F_{17}) - 4F_{23}Q_7 + 2(F_{23} - F_{24})(2r + Q_9 - 4S_2) + \\
& e^{\frac{1}{2}(\lambda-\nu)}[4F_{19}Q_6 + 2(F_{19} + F_{21})(-Q_8 + 4S_4) + \\
& 4F_3(Q_{10} - S_5) - 4F_9S_5] + 4F_7(Q_{11} - S_6) - 4F_{11} + 2r^2(F'_{14} - F'_{17}) = 0, \\
& 4rF_{13} + (F_{11} - F_7)(2r + Q_9 - 4S_2) - 2F_{11}Q_7 + \\
& e^{\frac{1}{2}(\lambda-\nu)}[(Q_8 - 4S_4)(F_9 - F_3) - 2F_9Q_6 - 2F_{21}(Q_{10} - S_5) - 2F_{19}S_5] - \\
& 2F_{24}(Q_{11} - S_6) + 2F_{23}S_6 + 2r^2F'_{13} = 0, \\
& 2e^{\frac{1}{2}(\lambda+\nu)}[(4S_4 - Q_8)(F_{18} + F_{20}) + 2Q_6F_{18} + 2F_2(Q_{10} - S_5) - \\
& 2F_8S_5 + r^2\dot{F}_{14}] - r^2(F_{15} + F_{16})(Q_1 + 4S_1 - e^\nu\nu') + 2r^2F_{15}Q_2 = 0, \\
& -r^2F_{15}(e^\lambda Q_0 + e^\nu Q_4) + 2e^{\frac{1}{2}(\lambda+3\nu)}[F_{20}(2r + Q_9 - 4S_2) - 2F_2(Q_{11} - S_6)] - \\
& 2e^{\lambda+\nu}[F_{22}(2Q_6 - Q_8 + 4S_4) + 2F_{10}S_5 - r^2\dot{F}_{15}] - \\
& e^{\frac{1}{2}(\lambda+\nu)}r^2(F_{14} + F_{17})(Q_1 + 4S_1 - e^\nu\nu') = 0,
\end{aligned}$$

$$\begin{aligned}
& r^2 F_{16}(e^\lambda Q_0 + e^\nu Q_4) + 2e^{\frac{1}{2}(\lambda+3\nu)}[F_{18}(2r - 2Q_7 + Q_9 - 4S_2) + F_8 S_6] + \\
& 2e^{\lambda+\nu}[-F_{25}(Q_8 - 4S_4) + 2F_6(Q_{10} - S_5) + r^2 \dot{F}_{16}] - \\
& e^{\frac{1}{2}(\lambda+\nu)} r^2 (F_{14} + F_{17})(Q_1 - 2Q_2 + 4S_1 - e^\nu \nu') = 0, \\
& e^\nu [4F_6(S_6 - Q_{11}) + 2(F_{25} - F_{22})(2r + Q_9 - 4S_2) + 4F_{22}Q_7 + \\
& 4F_{10}S_6 + r^2 \nu' (F_{15} + F_{16})] + r^2 [2F_{15}Q_2 - \\
& (Q_1 + 4S_1)(F_{15} + F_{16})] + 2e^{\frac{1}{2}(\lambda+\nu)} r^2 \dot{F}_{17} = 0, \\
& e^{\frac{1}{2}\lambda} [2F_2(Q_{10} - S_5) + F_{18}(2Q_6 - Q_8 + 4S_4) + F_{20}(4S_4 - Q_8) - \\
& 2F_8 S_5 + r^2 (\dot{F}_{14} - \dot{F}_{17})] + e^{\frac{1}{2}\nu} [2F_6(Q_{11} - S_6) + \\
& (F_{22} - F_{25})(2r + Q_9 - 4S_2) - 2Q_7 F_{22} - 2F_{10}S_6] = 0, \\
& e^{\frac{1}{2}(\nu-\lambda)} [(F_6 - F_{10})(2r + Q_9 - 4S_2) + 2F_{10}Q_7 + 2F_{25}(Q_{11} - S_6) - 2F_{22}S_6] + \\
& (F_2 - F_8)(Q_8 - 4S_4) + 2F_8 Q_6 + 2F_{20}(Q_{10} - S_5) + 2F_{18}S_5 - 2r^2 \dot{F}_{13} = 0, \\
& e^{\frac{1}{2}(\lambda+3\nu)} [F_2(-Q_9 + 2r + r^2 \nu') + F_1(2r - 2Q_7 + Q_9 - 4S_2) - 4F_{20}(Q_{11} - \\
& S_6 - 2S_8) + 4r^2 F_2'] + 2e^{\frac{1}{2}(\lambda+\nu)} r^2 [F_7(Q_1 + 4S_1 - e^\nu \nu') - F_2 Q_1] + \\
& 2e^\lambda r^2 F_3 Q_0 + 2e^{\lambda+\nu} [F_3(Q_8 - r^2 \dot{\lambda}) + (F_0 + F_{12})(2Q_6 - Q_8 + 4S_4) - 2F_{26}S_5 + \\
& 2F_{21}(Q_{10} - S_5 - 2S_7) + r^2 (-2\dot{F}_3 + F_6 \dot{\lambda})] - 2e^\nu r^2 F_6(2Q_3 - Q_4 + 4S_3) = 0, \\
& \frac{1}{r} e^{\frac{1}{2}\nu} [F_{20}(-2r + Q_9) + F_1 S_6 - 2F_2(Q_{11} - S_6 + 2S_8) - r^2 (2F_{20}' + \\
& F_{20} \nu')] - e^{\frac{1}{2}\lambda-\nu} r F_{21} Q_0 + e^{-\frac{1}{2}\nu} r [F_{20} Q_1 + F_{24}(-Q_1 + 4S_1 + e^\nu \nu')] + \\
& \frac{1}{r} e^{\frac{1}{2}\lambda} [-F_{21} Q_8 + F_{26}(-2Q_6 + Q_8 - 4S_4) - 2S_5(F_0 + F_{12}) + 2F_3(Q_{10} - \\
& S_5 - 2S_7) + r^2 (2\dot{F}_{21} + F_{21} \dot{\lambda})] - e^{-\frac{1}{2}\lambda} r F_{25}(-2Q_3 + Q_4 - 4S_3 + e^\lambda \dot{\lambda}) = 0, \\
& e^{\lambda+\nu} [(F_{12} - F_5)(2r - 2Q_7 + Q_9 - 4S_2) + 2F_{26}S_6 - 2F_{25}(Q_{11} - S_6 - 2S_8) + \\
& 2r^2 F_6' + F_6(-Q_9 + 2r + r^2 \nu')] + e^{\frac{1}{2}(\lambda+\nu)} r^2 [Q_4(F_2 - F_7) - 4F_2 S_3] + \\
& e^{\frac{1}{2}(3\lambda+\nu)} [2F_4(Q_6 + 2S_4) + Q_8(F_7 - F_4) + 2F_{24}(Q_{10} - S_5 - 2S_7) + r^2 (-2\dot{F}_7 + \\
& \dot{\lambda} F_2 - \dot{\lambda} F_7)] + e^\lambda r^2 F_3(Q_1 - 2Q_2 + 4S_1 - e^\nu \nu') + e^\nu r^2 F_6 Q_5 = 0,
\end{aligned} \tag{12}$$

$$\begin{aligned}
& -e^{\lambda+\nu}[F_{26}(2r-2Q_7+Q_9-4S_2)-2S_6(F_{12}-F_5)+2F_6(Q_{11}-S_6- \\
& 2S_8)+2r^2F'_{25}+F_{25}(-Q_9+2r+r^2\nu')]-e^{\frac{1}{2}(\lambda+\nu)}r^2[Q_4(F_{20}-F_{24})- \\
& 4F_{20}S_3]-e^{\frac{1}{2}(3\lambda+\nu)}[F_{24}Q_8+2F_4S_5-2F_7(Q_{10}-S_5-2S_7)+r^2(-2\dot{F}_{24}+ \\
& \dot{\lambda}F_{20}-\dot{\lambda}F_{24})]-e^\lambda r^2F_{21}(Q_1-2Q_2+4S_1-e^\nu\nu')-e^\nu r^2F_{25}Q_5=0, \\
& e^{\frac{1}{2}\nu}[-2F_8-2rF'_8-r\nu'(F_8-F_{11})]+e^{-\frac{1}{2}\nu}r[-F_8Q_1+F_{11}(-Q_1+2Q_2- \\
& 4S_1)]+e^{\frac{1}{2}\lambda-\nu}rF_9Q_0+e^{\frac{1}{2}\lambda}r[2\dot{F}_9+\dot{\lambda}(F_9-F_{10})]-e^{-\frac{1}{2}\lambda}rF_{10}(Q_4-4S_3)+ \\
& \frac{1}{r}e^{\frac{1}{2}\lambda}[F_9Q_8+(F_0+F_{12})(Q_8-4S_4)+2(F_{26}-F_{19})(Q_{10}-S_5)+4F_{19}S_7]+ \\
& \frac{1}{r}e^{\frac{1}{2}\nu}[-F_8Q_9+F_4(2r+Q_9-4S_2)+2F_{18}(Q_{11}-S_6-2S_8)]=0, \\
& e^{\frac{1}{2}\nu}[2F_{10}+2rF'_{10}+r\nu'(F_{10}-F_9)]+e^{-\frac{1}{2}\nu}rF_9(Q_1+4S_4)+ \\
& e^{-\frac{1}{2}\lambda}r[F_{11}Q_4+F_8(-2Q_3+Q_4-4S_3)]-e^{\frac{1}{2}\lambda}r[2\dot{F}_{11}+\dot{\lambda}(F_{11}-F_8)]- \\
& e^{-\lambda+\frac{1}{2}\nu}rF_{10}Q_5-\frac{1}{r}e^{\frac{1}{2}\lambda}[F_{11}Q_8+\frac{1}{2}F_1(Q_8-4S_4)-2F_{23}(Q_{10}- \\
& S_5-2S_7)]+\frac{1}{r}e^{\frac{1}{2}\nu}[F_{10}Q_9+(F_{12}-F_5)(2r+Q_9-4S_2)+ \\
& 2(F_{26}+F_{22})(Q_{11}-S_6)-4F_{22}S_8]=0, \\
& -e^{\frac{1}{2}\lambda-\nu}rF_{19}Q_0+e^{\frac{1}{2}\nu}[2F_{18}+2rF'_{18}+r\nu'(F_{18}+F_{23})]+e^{-\frac{1}{2}\nu}r[F_{18}Q_1+ \\
& F_{23}(-Q_1+Q_2-4S_1)]-e^{-\frac{1}{2}\lambda}rF_{22}(Q_4-4S_3)-e^{\frac{1}{2}\lambda}r[2\dot{F}_{19}+\dot{\lambda}(F_{19}+F_{22})]- \\
& \frac{1}{r}e^{\frac{1}{2}\lambda}[F_{19}Q_8+F_{26}(Q_8-4S_4)-2(F_0+F_{12}+F_9)(Q_{10}-S_5)+4F_9S_7]+ \\
& \frac{1}{r}e^{\frac{1}{2}\nu}[F_{18}Q_9+2(F_4-F_8)(-Q_{11}+S_6)-4F_8S_8]=0, \\
& -e^{\frac{1}{2}\nu-\lambda}rF_{22}Q_5+e^{\frac{1}{2}\nu}[2F_{22}+2rF'_{22}+r\nu'(F_{22}+F_{19})]-e^{-\frac{1}{2}\nu}rF_{19}(Q_1+4S_1)+ \\
& e^{-\frac{1}{2}\lambda}r[F_{23}Q_4-F_{18}(-2Q_3+Q_4-4S_3)]-e^{\frac{1}{2}\lambda}r[2\dot{F}_{23}+\dot{\lambda}(F_{23}+F_{18})]- \\
& \frac{1}{r}e^{\frac{1}{2}\lambda}[F_{23}Q_8-(F_1-2F_{11})(Q_{10}-S_5)-4F_{11}S_7]+\frac{1}{r}e^{\frac{1}{2}\nu}[F_{22}Q_9+ \\
& F_{26}(2r+Q_9-4S_2)+2(F_5-F_{12}-F_{10})(Q_{11}-S_6)+4F_{10}S_8]=0.
\end{aligned}$$

In the case of vanishing pseudoscalar functions S_i ($i = 5, \dots, 8$) and Q_i ($i = 10, 11$) the curvature tensor is determined by 14 functions \tilde{F}_i ($i = 0, 1, \dots, 13$) (other functions F_i ($i =$

14, ... 26) are equal to zero), expressions of which follow from (9):

$$\begin{aligned}
F^{\hat{0}\hat{0}}_{\hat{1}\hat{0}} &= \tilde{F}_0, \quad F^{\hat{0}\hat{1}}_{\hat{1}\hat{0}} = \tilde{F}_1, \quad F^{\hat{0}\hat{2}}_{\hat{2}\hat{0}} = F^{\hat{0}\hat{3}}_{\hat{3}\hat{0}} = \tilde{F}_2, \quad F^{\hat{0}\hat{2}}_{\hat{2}\hat{1}} = F^{\hat{0}\hat{3}}_{\hat{3}\hat{1}} = \tilde{F}_3, \\
F^{\hat{1}\hat{0}}_{\hat{1}\hat{0}} &= \tilde{F}_4, \quad F^{\hat{1}\hat{1}}_{\hat{1}\hat{0}} = \tilde{F}_5, \quad F^{\hat{1}\hat{2}}_{\hat{2}\hat{0}} = F^{\hat{1}\hat{3}}_{\hat{3}\hat{0}} = \tilde{F}_6, \quad F^{\hat{1}\hat{2}}_{\hat{2}\hat{1}} = F^{\hat{1}\hat{3}}_{\hat{3}\hat{1}} = \tilde{F}_7, \\
F^{\hat{2}\hat{0}}_{\hat{2}\hat{0}} &= F^{\hat{3}\hat{0}}_{\hat{3}\hat{0}} = \tilde{F}_8, \quad F^{\hat{2}\hat{0}}_{\hat{2}\hat{1}} = F^{\hat{3}\hat{0}}_{\hat{3}\hat{1}} = \tilde{F}_9, \quad F^{\hat{2}\hat{1}}_{\hat{2}\hat{0}} = F^{\hat{3}\hat{1}}_{\hat{3}\hat{0}} = \tilde{F}_{10}, \\
F^{\hat{2}\hat{1}}_{\hat{2}\hat{1}} &= F^{\hat{3}\hat{1}}_{\hat{3}\hat{1}} = \tilde{F}_{11}, \quad F^{\hat{2}\hat{2}}_{\hat{0}\hat{1}} = F^{\hat{3}\hat{3}}_{\hat{0}\hat{1}} = \tilde{F}_{12}, \quad -F^{\hat{3}\hat{2}}_{\hat{3}\hat{2}} = F^{\hat{2}\hat{3}}_{\hat{3}\hat{2}} = \tilde{F}_{13},
\end{aligned} \tag{13}$$

where explicit form of functions \tilde{F}_i is:

$$\begin{aligned}
\tilde{F}_0 &= \frac{1}{2} \{ e^{-\frac{3}{2}(\lambda+\nu)} [-Q_3(4S_1 + Q_1) + Q_2(2Q_3 - Q_4 + 4S_3)] + \\
&e^{-\frac{1}{2}(\lambda+3\nu)} (-\dot{Q}_1 - Q_2\dot{\lambda} + Q_1\dot{\nu} + Q'_0 - Q_0\nu') + e^{-\frac{1}{2}(3\lambda+\nu)} Q_3\nu' \}, \\
\tilde{F}_1 &= \frac{1}{4} \{ e^{-\lambda-2\nu} [Q_1^2 + Q_0(-2Q_3 + Q_4 - 4S_3 + e^\lambda\dot{\lambda}) + 4Q_1S_1] + \\
&e^{-2\lambda-\nu} [Q_5(Q_1 + 4S_1 - e^\nu\nu') + Q_4(Q_4 - 2Q_3 - 4S_3)] + e^{-(\lambda+\nu)} [4\dot{Q}_3 - \\
&2\dot{Q}_4 + 8\dot{S}_3 + Q_1\lambda' + (\dot{\lambda} + \dot{\nu})(Q_4 - 4S_3 - 2Q_3) + Q_4\dot{\lambda} - 2Q'_1 - \\
&8S'_1 + 4S_1(\lambda' + \nu') + e^\lambda(-\dot{\lambda}^2 + \dot{\lambda}\dot{\nu} - 2\ddot{\lambda})] - e^{-\lambda}(-\lambda'\nu' + \nu'^2 + 2\nu'') \}, \\
\tilde{F}_2 &= -\frac{1}{4r^4} \{ e^{-(\lambda+\nu)} r^2(2r - 2Q_7 + Q_9 - 4S_2)(Q_1 + 4S_1 - e^\nu\nu') + \\
&e^{-\nu} [(-Q_8 + 4S_4 + 2Q_6)(Q_8 + r^2\dot{\nu}) + r^2(-4\dot{Q}_6 + 2\dot{Q}_8 - \\
&8\dot{S}_4)] + e^{-2\nu} r^2 Q_0(2Q_6 - Q_8 + 4S_4) \}, \\
\tilde{F}_3 &= \frac{1}{4r^2} \{ e^{-\frac{1}{2}(\lambda+\nu)} [-\frac{1}{r^2}(2r + Q_9)(2Q_6 - Q_8 + 4S_4) + 8S'_4 - 8S_4\nu' + 4Q'_6 - \\
&2Q'_8 - 4\nu'Q_6 + 2\nu'Q_8] + e^{-\frac{1}{2}(3\lambda+\nu)} (2r - 2Q_7 + Q_9 - 4S_2)(-2Q_3 + \\
&Q_4 - 4S_1 + e^\lambda\dot{\lambda}) + e^{-\frac{1}{2}(\lambda+3\nu)} (2Q_6 - Q_8 + 4S_4)(-Q_1 + e^\nu\nu') \}, \\
\tilde{F}_4 &= \frac{1}{4} \{ e^{-\lambda-2\nu} [Q_0(Q_4 - 4S_3 + e^\lambda\dot{\lambda}) + Q_1(Q_1 - 2Q_2 + 4S_1 - e^\nu\lambda' - \\
&2e^\nu\nu')] + e^{-2\lambda-\nu} [Q_4(Q_4 - 4S_3) + Q_5(Q_1 + 4S_1 - 2Q_2)] + e^{-\lambda-\nu} [2\dot{Q}_4 - \\
&8\dot{S}_3 + 4S_3(\dot{\lambda} + \dot{\nu}) - Q_4\dot{\nu} + 2Q'_1 - 4Q'_2 + 8S'_1 + (\lambda' + \nu')(2Q_2 - 4S_1)] + \\
&e^{-\nu}(\dot{\lambda}^2 - \dot{\lambda}\dot{\nu} + 2\ddot{\lambda}) - e^{-2\lambda}Q_5\nu' + e^{-\lambda}(\lambda'\nu' - \nu'^2 - 2\nu'') \},
\end{aligned} \tag{14}$$

$$\begin{aligned}
\tilde{F}_5 &= \frac{1}{2}e^{-\frac{3}{2}(\lambda+\nu)}[-Q_3(Q_1+4S_1)+e^\nu(-\dot{Q}_5+Q_5\dot{\lambda}+Q'_4- \\
&\quad Q_4\lambda'+Q_3\nu')+Q_2(2Q_3-Q_4+4S_3-e^\lambda\dot{\lambda})], \\
\tilde{F}_6 &= -\frac{1}{4r^4}\{r^2e^{-\frac{1}{2}(\lambda+3\nu)}(2Q_6-Q_8+4S_4)(Q_1+4S_1- \\
&\quad e^\nu\nu'-2Q_2)+e^{-\frac{1}{2}(\lambda+\nu)}[(2r-2Q_7+Q_9-4S_2)(Q_8+r^2\dot{\lambda}- \\
&\quad e^{-\lambda}r^2Q_4)+r^2(4\dot{Q}_7-2\dot{Q}_9+8\dot{S}_2)]\}, \\
\tilde{F}_7 &= -\frac{1}{4r^4}\{-e^{-2\lambda}r^2(Q_5-e^\lambda\lambda')(2r-2Q_7+Q_9-4S_2)+ \\
&\quad e^{-\lambda-\nu}r^2(2Q_6-Q_8+4S_4)(4S_3-Q_4-e^\lambda\dot{\lambda})+e^{-\lambda}[Q_9(4r+Q_9- \\
&\quad 4S_2)-8rS_2+r^2(4Q'_7-2Q'_9+8S'_2)-2Q_7(2r+Q_9)]\}, \\
\tilde{F}_8 &= \frac{1}{4r^4}\{e^{-\nu}[(Q_8+e^{-\nu}r^2Q_0-r^2\dot{\nu})(Q_8-4S_4)+2r^2(\dot{Q}_8-4\dot{S}_4)]+ \\
&\quad e^{-\lambda-\nu}r^2(2r+Q_9-4S_2)(Q_1-2Q_2+4S_1-e^\nu\nu')\}, \\
\tilde{F}_9 &= \frac{1}{4r^4}e^{-\frac{1}{2}(\lambda+\nu)}[(Q_8-4S_4)(Q_9-2r+e^{-\nu}r^2Q_1-r^2\nu')- \\
&\quad e^{-\lambda}r^2(2r+Q_9-4S_2)(Q_4-4S_3+e^\lambda\dot{\lambda})+2r^2(Q'_8-4S'_4)], \\
\tilde{F}_{10} &= -\frac{1}{4r^4}e^{-\frac{1}{2}(\lambda+\nu)}[(2r+Q_9-4S_2)(Q_8-e^{-\lambda}r^2Q_4-r^2\dot{\lambda})+ \\
&\quad 2r^2(\dot{Q}_9-4\dot{S}_2)+e^{-\nu}r^2(Q_8-4S_4)(Q_1+4S_1-e^\nu\nu')], \\
\tilde{F}_{11} &= -\frac{1}{4r^4}e^{-\lambda}\{r^2[4+e^{-\nu}(Q_8-4S_4)(2Q_3-Q_4+4S_3-e^\lambda\dot{\lambda})+ \\
&\quad 2Q'_9-8S'_2]+(2r+Q_9-4S_2)(Q_9-2r-e^{-\lambda}r^2Q_5+r^2\lambda')\}, \\
\tilde{F}_{12} &= \frac{1}{2r^3}e^{-\frac{1}{2}(\lambda+\nu)}[2Q_8+r(\dot{Q}_9-Q'_8)], \\
\tilde{F}_{13} &= -\frac{1}{4r^4}[4r^2-e^{-\lambda}(2r+Q_9-4S_2)(2r-2Q_7+Q_9-4S_2)+ \\
&\quad e^{-\nu}(Q_8-4S_4)(-2Q_6+Q_8-4S_4)].
\end{aligned}$$

In this case Bianchi identities are reduced to 6 following relations:

$$\begin{aligned}
&-4r\tilde{F}_{13}+(2r+Q_9-4S_2)(\tilde{F}_7-\tilde{F}_{11})+2\tilde{F}_{11}Q_7+ \\
&e^{\frac{1}{2}(\lambda-\nu)}[(Q_8-4S_4)(\tilde{F}_3-\tilde{F}_9)+2\tilde{F}_9Q_6]-2r^2\tilde{F}'_{13}=0,
\end{aligned}$$

$$\begin{aligned}
& e^{\frac{1}{2}(\nu-\lambda)}[(\tilde{F}_{10} - \tilde{F}_6)(2r + Q_9 - 4S_2) - 2\tilde{F}_{10}Q_7] + \\
& (Q_8 - 4S_4)(\tilde{F}_8 - \tilde{F}_2) - 2\tilde{F}_8Q_6 + 2r^2\dot{\tilde{F}}_{13} = 0, \\
& e^{\frac{1}{2}\nu}[\tilde{F}_2(\frac{1}{r}Q_9 - 2 - r\nu') + \frac{1}{r}\tilde{F}_1(2r - 2Q_7 + Q_9 - 4S_2) - 2r\tilde{F}'_2] - e^{\frac{1}{2}\lambda-\nu}r\tilde{F}_3Q_0 + \\
& re^{-\frac{1}{2}\nu}[\tilde{F}_2Q_1 + \tilde{F}_7(-Q_1 - 4S_1 + e^\nu\nu')] + e^{\frac{1}{2}\lambda}\{\frac{1}{r}[-\tilde{F}_3Q_8 + (\tilde{F}_{12} - \tilde{F}_0)(2Q_6 - \\
& Q_8 + 4S_4)] + r(2\dot{\tilde{F}}_3 + \tilde{F}_3\dot{\lambda})\} - e^{-\frac{1}{2}\lambda}r\tilde{F}_6(-2Q_3 + Q_4 - 4S_3 + e^\lambda\dot{\lambda}) = 0, \\
& e^{\frac{1}{2}\nu}\{\tilde{F}_6(-2 - r\nu') - 2r\tilde{F}'_6 + \frac{1}{r}[\tilde{F}_6Q_9 + (\tilde{F}_{12} + \tilde{F}_5)(2r - 2Q_7 + Q_9 - 4S_2)]\} + \\
& e^{-\frac{1}{2}\nu}r\tilde{F}_3(-Q_1 + 2Q_2 - 4S_1 + e^\nu\nu') + e^{-\frac{1}{2}\lambda}r[\tilde{F}_7Q_4 - \tilde{F}_2(Q_4 - 4S_3 + e^\lambda\dot{\lambda})] - \\
& e^{-\lambda+\frac{1}{2}\nu}r\tilde{F}_6Q_5 + e^{\frac{1}{2}\lambda}\{\frac{1}{r}[-\tilde{F}_7Q_8 - \tilde{F}_4(2Q_6 - Q_8 + 4S_4)] + 2r\dot{\tilde{F}}_7 + r\tilde{F}_7\dot{\lambda}\} = 0, \quad (15) \\
& e^{\frac{1}{2}\nu}\{2\tilde{F}_8 + \frac{1}{r}[\tilde{F}_8Q_9 - \tilde{F}_4(2r + Q_9 - 4S_2)] + 2r\tilde{F}'_8 + r\nu'(\tilde{F}_8 - \tilde{F}_{11})\} + \\
& e^{-\frac{1}{2}\nu}r[\tilde{F}_8Q_1 + \tilde{F}_{11}(Q_1 - 2Q_2 + 4S_1)] - e^{\frac{1}{2}\lambda-\nu}r\tilde{F}_9Q_0 + e^{\frac{1}{2}\lambda}\{\frac{1}{r}[(\tilde{F}_{12} - \\
& \tilde{F}_0)(Q_8 - 4S_4) - \tilde{F}_9Q_8] - 2r\dot{\tilde{F}}_9 + r\dot{\lambda}(\tilde{F}_{10} - \tilde{F}_9)] + e^{-\frac{1}{2}\lambda}r\tilde{F}_{10}(Q_4 - 4S_3) = 0, \\
& e^{\frac{1}{2}\nu}\{\frac{1}{r}[(\tilde{F}_{12} + \tilde{F}_5)(2r + Q_9 - 4S_2) - \tilde{F}_{10}Q_9] - 2\tilde{F}_{10} - 2r\tilde{F}'_{10} + r\nu'(\tilde{F}_9 - \tilde{F}_{10})\} - \\
& e^{-\frac{1}{2}\nu}r\tilde{F}_9(Q_1 + 4S_1) + e^{-\lambda+\frac{1}{2}\nu}r\tilde{F}_{10}Q_5 + e^{\frac{1}{2}\lambda}\{\frac{1}{r}[\tilde{F}_{11}Q_8 + \tilde{F}_1(Q_8 - 4S_4)] + \\
& 2r\dot{\tilde{F}}_{11} + r\dot{\lambda}(\tilde{F}_{11} - \tilde{F}_8)\} - e^{-\frac{1}{2}\lambda}r[\tilde{F}_{11}Q_4 + \tilde{F}_8(Q_4 - 2Q_3 - 4S_3)] = 0.
\end{aligned}$$

Now let us consider spherically-symmetric gravitational fields in the frame of MAGT corresponding to particular gravitational Lagrangian:

$$L_G = f_0 F + f F^2 + a S_{\nu\mu\alpha} S^{\alpha\mu\nu} + k Q_{\mu\nu\lambda} Q^{\mu\lambda\nu} + m Q^\alpha{}_{\lambda\alpha} S_\beta{}^{\beta\lambda}, \quad (16)$$

where $f_0 = (16\pi G)^{-1}$, G is Neuton's gravitational constant; f, a, k, m are indefinite parameters, $F = F^{\mu\nu}{}_{\mu\nu}$.

The gravitational equations can be obtained by variation of total action integral

$$I = \int \delta^4 x h (L_G + L_m) \quad (17)$$

(L_m is Lagrangian of matter and $h = \det(h^i{}_\mu)$) with respect to $h^i{}_\mu$ and $A^{ik}{}_\mu$. As result of variation we get 16 h-equations:

$$H_i{}^\mu - \nabla_\nu \sigma_i{}^{\mu\nu} = t_i{}^\mu, \quad (18)$$

and 64 A-equations:

$$2\nabla_\nu \varphi_{ik}{}^{\nu\mu} + \sigma_{ik}{}^\mu = -J_{ik}{}^\mu, \quad (19)$$

were $H_i^\mu = h^{-1}(\delta L_G / \delta h^i_\mu)$, $\sigma_i^{\mu\nu} = (\partial L_G / \partial S^i_{\mu\nu})$, $\varphi_{ik}^{\mu\nu} = (\partial L_G / \partial F^{ik}_{\mu\nu})$, $t_i^\mu = -h^{-1}(\delta L_m / \delta h^i_\mu)$, $J_{ik}^\mu = -h^{-1}(\delta L_m / \delta A^{ik}_\mu)$.

In spherically-symmetric case with vanishing pseudoscalar torsion and nonmetricity functions the system of gravitational equations (18) – (19) is reduced to 19 differential equations:

$$\begin{aligned}
& 2f_0(\tilde{F}_{19} - \tilde{F}_{20} - \tilde{F}_{17}) + f[4\tilde{F}_{19}^2 - \tilde{F}_{14}^2 - (2\tilde{F}_1 - \tilde{F}_4)^2 + 4(\tilde{F}_{20} + \tilde{F}_{17})^2 - \\
& 8\tilde{F}_{19}(\tilde{F}_{20} + \tilde{F}_{17}) + 4\tilde{F}_8(2\tilde{F}_1 - \tilde{F}_4) - 4\tilde{F}_8^2 + 2\tilde{F}_{14}(\tilde{F}_4 - 2\tilde{F}_1 + 2\tilde{F}_8)] + \\
& k\{e^{-\lambda-2\nu}Q_2^2 - e^{-3\nu}Q_0^2 + e^{-2\lambda-\nu}Q_3^2 - e^{-3\lambda}Q_5^2 + \frac{1}{r^4}[2e^{-\nu}Q_6^2 - \\
& 2e^{-\lambda}Q_7(Q_7 + 2Q_9)]\} + \frac{m}{4}\{e^{-\lambda-2\nu}(Q_1Q_2 + 4Q_2S_1 - Q_0Q_4) + \\
& \frac{4}{r}(e^{-\lambda} - \nu Q_2 - e^{-2\lambda}Q_5) + e^{-2\lambda-\nu}(Q_3Q_4 - Q_1Q_5 - 4Q_3S_3) + \\
& \frac{2}{r^2}[e^{-\lambda-\nu}(Q_4Q_6 - 4Q_6S_3 - Q_1Q_7 + Q_3Q_8 - 4Q_3S_4) - \\
& e^{-2\nu}Q_0Q_8 + 4e^{-2\lambda}Q_5S_2 + e^{-\nu}Q_6\dot{\lambda} - 2e^{-\lambda}Q_7' + e^{-\lambda}Q_7\lambda'] + \\
& \frac{4}{r^4}[e^{-\nu}Q_6(Q_8 - 4S_4) + 4e^{-\lambda}Q_7S_2] + e^{-2\nu}Q_0\dot{\lambda} + \\
& e^{-2\lambda}(3Q_5\lambda' - 2Q_5') + e^{-\lambda-\nu}(Q_3\dot{\lambda} + 2Q_2' - Q_2\lambda' - 2Q_2\nu')\} + \\
& \frac{a}{2}\{e^{-\lambda-\nu}(\frac{4}{r}S_1 - S_3\dot{\lambda} + 2S_1' - S_1\lambda' - 2S_1\nu') + e^{-\lambda-2\nu}(Q_1S_1 + \\
& 2S_1^2) - \frac{1}{r^4}[4e^{-\lambda}S_2^2 + 2e^{-\nu}(Q_8S_4 - 2S_4^2)] + e^{-2\lambda-\nu}(2S_3^2 - Q_4S_3)\} = t_0^{\hat{0}}, \\
& (\tilde{F}_5 - \tilde{F}_{18})[2f_0 + 4f(2\tilde{F}_{19} - \tilde{F}_{14} - 2\tilde{F}_{20} - 2\tilde{F}_1 + \tilde{F}_4 - 2\tilde{F}_{17} + \\
& 2\tilde{F}_8)] + 2k[e^{-\frac{1}{2}(\lambda+5\nu)}Q_0Q_2 - e^{-\frac{3}{2}(\lambda+\nu)}Q_2(Q_3 + Q_4) + \\
& e^{-\frac{1}{2}(5\lambda+\nu)}Q_4Q_5 + \frac{2}{r^4}e^{-\frac{1}{2}(\lambda+\nu)}Q_7Q_8] + \frac{m}{4}\{e^{-\frac{1}{2}(\lambda+5\nu)}Q_0(Q_2 - \\
& Q_1 - 4S_1) + e^{-\frac{3}{2}(\lambda+\nu)}(Q_3Q_1 - 2Q_3Q_2 + Q_0Q_5 + 4S_1Q_3 - \\
& 4S_1Q_4 + 4Q_2S_3) + \frac{1}{r^2}e^{-\frac{1}{2}(\lambda+3\nu)}[2Q_6(Q_1 - 2Q_2 + 4S_1) + \\
& 2Q_0Q_7 + 2Q_2Q_8 + 8Q_2S_4 + r^2(Q_2\dot{\lambda} - 2\dot{Q}_2 + 2Q_2\dot{\nu} + Q_0\nu')]\} + \\
& e^{-\frac{1}{2}(3\lambda+\nu)}[2\dot{Q}_5 - 3Q_5\dot{\lambda} - \frac{1}{r^2}(2Q_5Q_8 + Q_4S_2) - Q_3\nu'] + \\
& \frac{1}{r^2}e^{-\frac{1}{2}(\lambda+\nu)}(4\dot{Q}_7 - \frac{4}{r^2}Q_7Q_8 - 2Q_7\dot{\lambda} - Q_6\nu')\} + \\
& \frac{a}{2}[\frac{2}{r^4}e^{-\frac{1}{2}(\lambda+\nu)}Q_8S_2 - e^{-\frac{1}{2}(\lambda+5\nu)}Q_0S_1 + e^{-\frac{3}{2}(\lambda+\nu)}S_3(2Q_2 - Q_1 - 4S_1) + \\
& e^{-\frac{1}{2}(\lambda+3\nu)}(S_1\dot{\lambda} - 2\dot{S}_1 + 2S_1\dot{\nu}) + e^{-\frac{1}{2}(3\lambda+\nu)}S_3\nu'] = t_1^{\hat{0}},
\end{aligned}$$

$$\begin{aligned}
& 2(\tilde{F}_{15} - \tilde{F}_9)[f_0 + 2f(2\tilde{F}_{19} - \tilde{F}_{14} - 2\tilde{F}_{20} - 2\tilde{F}_1 + 2\tilde{F}_4 - 2\tilde{F}_{17} + \\
& 2\tilde{F}_8)] + 2k[e^{-\frac{3}{2}(\lambda+\nu)}Q_3(O_1 + Q_2) - e^{-\frac{1}{2}(\lambda+5\nu)}Q_0Q_1 - \\
& e^{-\frac{1}{2}(5\lambda+\nu)}Q_3Q_5 - \frac{2}{r^4}e^{-\frac{1}{2}(\lambda+\nu)}Q_6Q_9] + \frac{m}{4}\{e^{-\frac{3}{2}(\lambda+\nu)}(2Q_2Q_3 - \\
& Q_2Q_4 - Q_0Q_5 + 4Q_3S_1 - 4Q_1S_3 + 4Q_2S_3) + e^{-\frac{1}{2}(5\lambda+\nu)}Q_5(Q_4 - Q_3 - 4S_3) + \\
& e^{-\frac{1}{2}(\lambda+\nu)}[\frac{8}{r^3}Q_6 + \frac{4}{r^4}Q_6Q_9 + \frac{2}{r^2}(Q_7\dot{\lambda} - 2Q'_6 + Q_6\nu')]\} + e^{-\frac{1}{2}(3\lambda+\nu)}[\frac{2}{r^2}(Q_5Q_6 - \\
& 2Q_3Q_7 + Q_4Q_7 + Q_3Q_9 + 4Q_3S_2 - 4Q_7S_3) + Q_5\dot{\lambda} - 2Q'_3 + 2Q_3\lambda' + Q_3\nu'] + \\
& e^{-\frac{1}{2}(\lambda+3\nu)}[2Q'_0 - 3Q_0\nu' - \frac{2}{r^2}(Q_0Q_9 + 4Q_1S_4) - Q_2\dot{\lambda}]\} + \frac{a}{2}[e^{-\frac{3}{2}(\lambda+\nu)}S_1(2Q_3 - \\
& Q_4 + 4S_3) + e^{-\frac{1}{2}(3\lambda+\nu)}(\frac{4}{r}S_3 + 2S'_3 + 2S_3\lambda' - S_3\nu') + e^{-\frac{1}{2}(\lambda+\nu)}Q_5S_3 - \\
& e^{-\frac{1}{2}(\lambda+\nu)}(\frac{4}{r^3}S_4 + \frac{2}{r^4}Q_9S_4) - e^{-\frac{1}{2}(\lambda+3\nu)}S_1\dot{\lambda}] = t_0^{\hat{1}}, \\
& 2f_0(\tilde{F}_8 - \tilde{F}_1 - \tilde{F}_{20}) + f[8\tilde{F}_{20}\tilde{F}_1 - \tilde{F}_{14}^2 - 4\tilde{F}_{19}^2 + 4\tilde{F}_1^2 - 4\tilde{F}_{19}(\tilde{F}_4 - 2\tilde{F}_{17}) + \\
& 2\tilde{F}_{14}(2\tilde{F}_{19} + \tilde{F}_4 - 2\tilde{F}_7) - (\tilde{F}_4 - 2\tilde{F}_{17})^2 - 8\tilde{F}_1\tilde{F}_8 + 4(\tilde{F}_{20} - \tilde{F}_8)^2] + \\
& k[e^{-3\nu}Q_0^2 - e^{-\lambda-2\nu}Q_2^2 - e^{-2\lambda-\nu}Q_3^2 + e^{-3\lambda}Q_5^2 + \frac{1}{r^4}e^{-\nu}(2Q_6^2 + 4Q_6Q_8) - \\
& \frac{2}{r^4}e^{-\lambda}Q_7^2] + m\{e^{-\lambda-\nu}[\frac{1}{r}Q_2 - \frac{1}{2r^2}(Q_4Q_6 - Q_1Q_7 - Q_2Q_9 - 4Q_7S_1 + \\
& 4Q_2S_2) + \frac{1}{2}\dot{Q}_3 - \frac{1}{2}Q_3\dot{\lambda} - \frac{1}{4}Q_3\dot{\nu} + \frac{1}{4}Q_2\nu'] + \\
& \frac{1}{4}e^{-\lambda-2\nu}(Q_0Q_4 - Q_1Q_2 - 4Q_2S_1) + \frac{1}{4}e^{-2\lambda-\nu}(Q_1Q_5 - Q_3Q_4 + \\
& 4Q_3S_3) - e^{-2\lambda}Q_5(\frac{1}{r} + \frac{1}{2r^2}Q_9 + \frac{1}{4}\nu') + e^{-\lambda}Q_7(\frac{4}{r^4}S_2 - \frac{2}{r^3} - \frac{1}{r^4}Q_9 - \frac{1}{2r^2}\nu') + \\
& e^{-2\nu}(\frac{2}{r^2}Q_0S_4 - \frac{1}{2}\dot{Q}_0 + \frac{3}{4}Q_0\dot{\nu}) + \frac{1}{r^2}e^{-\nu}(\dot{Q}_6 - \frac{4}{r^2}Q_6S_4 - \frac{1}{2}Q_6\dot{\nu})\} + a[\frac{1}{r^3}e^{-\lambda}S_2(2 + \\
& \frac{1}{r}Q_9 - \frac{2}{r}S_2) - e^{-\lambda-2\nu}(\frac{1}{2}Q_1S_1 + S_1^2) + e^{-2\lambda-\nu}(\frac{1}{2}Q_4S_3 - S_3^2) + \frac{2}{r^4}e^{-\nu}S_4^2 + \\
& e^{-\lambda-\nu}(S_3\dot{\lambda} - \dot{S}_3 + \frac{1}{2}S_3\dot{\nu} + \frac{1}{2}S_1\nu')] = t_1^{\hat{1}},
\end{aligned}$$

$$\begin{aligned}
& f_0(\tilde{F}_4 - \tilde{F}_1 + \tilde{F}_{19} - \tilde{F}_{17} + \tilde{F}_8 - \tilde{F}_{14}) + f[\tilde{F}_{14}^2 - 4\tilde{F}_{20}^2 - \\
& 4\tilde{F}_{20}\tilde{F}_1 - 2\tilde{F}_1\tilde{F}_4 + \tilde{F}_4^2 + (4\tilde{F}_{20} + 2\tilde{F}_4)(\tilde{F}_{19} - \tilde{F}_{17} + \tilde{F}_8) - \\
& 2\tilde{F}_{14}(\tilde{F}_{19} - \tilde{F}_1 + \tilde{F}_4 - \tilde{F}_{17} + \tilde{F}_8)] + k[e^{-3\nu}Q_0^2 - \\
& e^{-\lambda-2\nu}(2Q_1Q_2 + Q_2^2) + e^{-2\lambda-\nu}(Q_3^2 + 2Q_3Q_4) - e^{-3\lambda}Q_5^2 + \frac{2}{r^4}e^{-\nu}Q_6Q_8 - \\
& \frac{2}{r^4}e^{-\lambda}Q_7Q_9] + m\{e^{-\lambda-\nu}[\frac{1}{2r}Q_2 + \frac{1}{r^2}(Q_7S_1 - \frac{1}{4}Q_3Q_8 - \frac{1}{4}Q_2Q_9 - \\
& Q_2S_2 - Q_6S_3 - Q_3S_4) + \frac{1}{2}\dot{Q}_3 - \frac{1}{4}Q_3\dot{\lambda} - \frac{1}{4}Q_3\dot{\nu} + \frac{1}{2}Q_2' - \\
& \frac{1}{4}Q_2\lambda' - \frac{1}{4}Q_2\nu'] + e^{-2\lambda}[Q_5(\frac{1}{4r^2}Q_9 + \frac{1}{r^2}S_2 - \frac{1}{2r} + \\
& \frac{3}{4}\lambda' - \frac{1}{4}\nu') - \frac{1}{2}Q_5'] + e^{-\lambda}[Q_7(\frac{1}{r^3} + \frac{1}{2r^4}Q_9 + \\
& \frac{1}{2r^2}\lambda' - \frac{1}{2r^2}\nu') - \frac{1}{r^2}Q_7'] + e^{-2\nu}[Q_0(\frac{1}{4r^2}Q_8 + \frac{1}{r^2}S_4 - \frac{1}{4}\dot{\lambda} + \frac{3}{4}\dot{\nu}) - \\
& \frac{1}{2}\dot{Q}_0] + \frac{1}{2r^2}e^{-\nu}[2\dot{Q}_6 + Q_6(\dot{\lambda} - \dot{\nu} - \frac{1}{r^2}Q_8)] + e^{-\lambda-2\nu}(Q_0S_3 - Q_2S_1) + \\
& e^{-2\lambda-\nu}(Q_5S_1 - Q_3S_3)\} + a[\frac{1}{2r^2}e^{-\lambda}(2S_2' - S_2\lambda' + S_2\nu' - \frac{2}{r}S_2 - \frac{1}{r^2}Q_9S_2) - \\
& e^{-\lambda-2\nu}S_1^2 + e^{-2\lambda-\nu}S_3^2 + \frac{1}{2r^2}e^{-\nu}(\frac{1}{r^2}Q_8S_4 - 2\dot{S}_4 - S_4\dot{\lambda} + S_4\dot{\nu})] = t_2^{\hat{2}}, \\
& 4e^{-\frac{3}{2}\nu}kQ_0 + (e^{-\lambda-\frac{1}{2}\nu}Q_3 + \frac{2}{r^2}e^{-\frac{1}{2}\nu}Q_6)[f_0 + 2f(2F_{19} - F_{14} - 2F_{20} - \\
& 2F_1 + F_4 - 2F_{17} + 2F_8)] = J_{00}^0, \\
& -\frac{1}{2}e^{-\frac{1}{2}\lambda-\nu}Q_2[2f_0 + 4f(2F_{19} - F_{14} - 2F_{20} - 2F_1 + F_4 - 2F_{17} + 2F_8) - \\
& 1 + 8k] - \frac{1}{2}e^{-\frac{3}{2}\lambda}Q_5 - \frac{1}{r^2}e^{-\frac{1}{2}\lambda-\nu}(e^\nu Q_7 - ar^2S_1) = J_{00}^1, \\
& \frac{1}{2}e^{-\frac{1}{2}\lambda-\nu}[4k(Q_1 + Q_2) + 2aS_1] + [\frac{1}{2}e^{-\frac{1}{2}\lambda-\nu}Q_1 + \frac{1}{2}e^{-\frac{3}{2}\lambda}Q_5 + \frac{1}{r^2}e^{-\frac{1}{2}\lambda}(2Q_7 - Q_9 + \\
& 4S_2)][f_0 + 2f(2F_{19} - F_{14} - 2F_{20} - 2F_1 + F_4 - 2F_{17} + 2F_8)] + 2fe^{-\frac{1}{2}\lambda}(2F_{19}' - \\
& F_{14}' - 2F_{20}' - 2F_1' + F_4' - 2F_{17}' + 2F_8') = J_{01}^0, \\
& [\frac{1}{r^2}e^{-\frac{1}{2}\nu}(Q_8 - 2S_4) - \frac{1}{2}e^{-\frac{3}{2}\nu}Q_0 - \frac{1}{2}e^{-\lambda-\frac{1}{2}\nu}Q_4][f_0 + 2f(2F_{19} - F_{14} - 2F_{20} - \\
& 2F_1 + F_4 - 2F_{17} + 2F_8)] - 2e^{-\lambda-\frac{1}{2}\nu}k(Q_3 + Q_4) + \frac{1}{2}e^{-\lambda-\frac{1}{2}\nu}S_3 + \frac{1}{r^2}e^{-\frac{1}{2}\nu}S_4 - \\
& fe^{-\frac{1}{2}\nu}(2\dot{F}_{19} - \dot{F}_{14} - 2\dot{F}_{20} - 2\dot{F}_1 + \dot{F}_4 - 2\dot{F}_{17} + 2\dot{F}_8) = J_{01}^1, \\
& (\frac{1}{2}e^{-\lambda-\frac{1}{2}\nu}Q_4 - \frac{1}{2}e^{-\frac{3}{2}\nu}Q_0 - 2e^{-\lambda-\frac{1}{2}\nu}S_3 - \frac{2}{r^2}e^{-\frac{1}{2}\nu}S_4)[f_0 + 2f(2F_{19} - F_{14} - 2F_{20} - \\
& 2F_1 + F_4 - 2F_{17} + 2F_8)] - \frac{2}{r^2}e^{-\frac{1}{2}\nu}k(Q_6 + Q_8) + \frac{4}{r^2}e^{-\frac{1}{2}\nu}S_4 + 2e^{-\lambda-\frac{1}{2}\nu}S_3 - \\
& 2fe^{-\frac{1}{2}\nu}(2\dot{F}_{19} - \dot{F}_{14} - 2\dot{F}_{20} - 2\dot{F}_1 + \dot{F}_4 - 2\dot{F}_{17} + 2\dot{F}_8) = J_{02}^2,
\end{aligned}$$

$$\begin{aligned}
& (\frac{1}{2}e^{-\frac{1}{2}\lambda-\nu}Q_1 + \frac{1}{2}e^{-\frac{3}{2}\lambda}Q_5 + \frac{1}{r^2}e^{-\frac{1}{2}\lambda}Q_9 - \frac{4}{r^2}e^{-\frac{1}{2}\lambda}S_2)[f_0 + 2f(2F_{19} - F_{14} - 2F_{20} - \\
& 2F_1 + F_4 - 2F_{17} + 2F_8)] + 2e^{-\frac{1}{2}\lambda-\nu}k(Q_1 + Q_2) - 2fe^{-\frac{1}{2}\lambda}(2F'_{19} - F'_{14} - \\
& 2F'_{20} - 2F'_1 + F'_4 - 2F'_{17} + 2F'_8) = J_{10}^0, \\
& [\frac{1}{r^2}e^{-\frac{1}{2}\nu}(2Q_6 - Q_8 + 4S_4) - \frac{1}{2}e^{-\frac{3}{2}\nu}Q_0 - \frac{1}{2}e^{-\lambda-\frac{1}{2}\nu}Q_4][f_0 + \\
& 2f(2F_{19} - F_{14} - 2F_{20} - 2F_1 + F_4 - 2F_{17} + 2F_8)] + \\
& 2fe^{-\frac{1}{2}\nu}(\dot{F}_{19} - \dot{F}_{14} - 2\dot{F}_{20} - 2\dot{F}_1 + \dot{F}_4 - 2\dot{F}_{17} + 2\dot{F}_8) + \frac{1}{2}e^{-\frac{3}{2}\nu}Q_0 - \\
& \frac{1}{2}e^{-\lambda-\frac{1}{2}\nu}[Q_3(1 + 4k) + 4kQ_4 - 2S_3(1 + a)] + \frac{1}{r^2}e^{-\frac{1}{2}\nu}(2S_4 - Q_6) = J_{10}^1, \\
& e^{-\lambda-\frac{1}{2}\nu}Q_3[f_0 + 2f(2F_{19} - F_{14} - 2F_{20} - 2F_1 + F_4 - 2F_{17} + 2F_8)] + \\
& e^{-\lambda-\frac{1}{2}\nu}(4kQ_3 + aS_3) = J_{11}^0, \\
& (\frac{2}{r^2}e^{-\frac{1}{2}\lambda}Q_7 - e^{-\frac{1}{2}\lambda-\nu}Q_2)[f_0 + 2f(2F_{19} - F_{14} - 2F_{20} - 2F_1 + F_4 - \\
& 2F_{17} + 2F_8)] - 4e^{-\frac{3}{2}\lambda}kQ_5 - 2e^{-\frac{1}{2}\lambda-\nu}S_1 - \frac{4}{r^2}e^{-\frac{1}{2}\lambda}S_2 = J_{11}^1, \\
& [\frac{1}{2}e^{-\frac{3}{2}\lambda}Q_5 - \frac{1}{2}e^{-\frac{1}{2}\lambda-\nu}(Q_1 + 4S_1) - \frac{2}{r^2}e^{-\frac{1}{2}\lambda}S_2][f_0 + 2f(2F_{19} - F_{14} - \\
& 2F_{20} - 2F_1 + F_4 - 2F_{17} + 2F_8)] - 2fe^{-\frac{1}{2}\lambda}(2F'_{19} - F'_{14} - 2F'_{20} - 2F'_1 + \\
& F'_4 - 2F'_{17} + 2F'_8) + \frac{2}{r^2}e^{-\frac{1}{2}\lambda}[2S_2 - k(Q_7 + Q_9)] + 2e^{-\frac{1}{2}\lambda-\nu}S_1 = J_{12}^2, \\
& [\frac{1}{2}e^{-\lambda-\frac{1}{2}\nu}(2Q_3 - Q_4 + 4S_3) - \frac{1}{2}e^{-\frac{3}{2}\nu}Q_0 + \frac{1}{r^2}e^{-\frac{1}{2}\nu}(Q_6 - Q_8 + 2S_4)][f_0 + \\
& 2f(2F_{19} - F_{14} - 2F_{20} - 2F_1 + F_4 - 2F_{17} + 2F_8)] + 2fe^{-\frac{1}{2}\nu}(\dot{F}_{19} - \\
& \dot{F}_{14} - 2\dot{F}_{20} - 2\dot{F}_1 + \dot{F}_4 - 2\dot{F}_{17} + 2\dot{F}_8) + e^{-\frac{3}{2}\nu}Q_0 + e^{-\lambda-\frac{1}{2}\nu}(2S_3 - Q_3) + \\
& \frac{1}{r^2}e^{-\frac{1}{2}\nu}[2Q_8 - 2Q_6(1 + k) + S_4(1 + a)] = J_{20}^2, \\
& [f_0 + 2f(2F_{19} - F_{14} - 2F_{20} - 2F_1 + F_4 - 2F_{17} + 2F_8)] \\
& [e^{-\frac{1}{2}\lambda-\nu}(\frac{1}{2}Q_1 - Q_2 + 2S_1) + \frac{1}{2}e^{-\frac{3}{2}\lambda}Q_5 + \frac{1}{r^2}e^{-\frac{1}{2}\lambda}(Q_7 - \\
& Q_9 + 2S_2)] + 2fe^{-\frac{1}{2}\lambda}(2F'_{19} - F'_{14} - 2F'_{20} - 2F'_1 + F'_4 - \\
& 2F'_{17} + 2F'_8) + e^{-\frac{1}{2}\lambda-\nu}(Q_2 + 2S_1) - e^{-\frac{3}{2}\lambda}Q_5 - \\
& \frac{1}{r^2}e^{-\frac{1}{2}\lambda}[2Q_7(1 + k) + 2kQ_9 - S_2(4 + a)] = J_{21}^2, \\
& \frac{1}{r^2}e^{-\frac{1}{2}\nu}(4kQ_6 + aS_4) + \frac{1}{r^2}e^{-\frac{1}{2}\nu}Q_6[f_0 + 2f(2F_{19} - \\
& F_{14} - 2F_{20} - 2F_1 + F_4 - 2F_{17} + 2F_8)] = J_{22}^0,
\end{aligned} \tag{20}$$

$$\begin{aligned} & \frac{1}{2}e^{-\frac{3}{2}\lambda}Q_5 - \frac{1}{2}e^{-\frac{1}{2}\lambda-\nu}Q_2 - \frac{1}{r^2}e^{-\frac{1}{2}\lambda}[Q_7(4k-1) + aS_2] - \\ & \frac{1}{r^2}e^{-\frac{1}{2}\lambda}Q_7[f_0 + 2f(2F_{19} - F_{14} - 2F_{20} - 2F_1 + F_4 - 2F_{17} + 2F_8)] = J_{22}{}^1. \end{aligned}$$

In the case $t_i{}^\mu = 0$, $J_{ik}{}^\mu = 0$ this system of equations is satisfied by vanishing torsion and nonmetricity and vacuum Schwarzschild metrics:

$$g_{\mu\nu} = \text{diag}((1 - r_g/r), -(1 - r_g/r)^{-1}, -r^2, -r^2 \sin^2 \theta), (r_g = \text{const}). \quad (21)$$

References

- [1] Hehl F. W., McCrea G. D., Mielke E. W. and Ne'eman Y. Metric-Affine Gauge Theory of Gravity: Fields Equations, Noether Identities, World Spinors, and Breaking of Dilaton Invariance. 1995, *Phys. Rep.* **258**, 1.
- [2] Minkevich A. V., Garkun A. S. Homogeneous isotropic models in the metric-affine gauge theory of gravity. 2000, *Class. Quantum Grav.* **17**, p.3045-3054.
- [3] Minkevich A. V. Spherically-symmetric spaces in the metric-affine gauge theory of gravitation, 1995, *Vestnik BGU, ser. 1*, No 3, pp. 30-33.